


AD-A156 754	ESTIMATION BY MOMENTS IN A MODEL OF FAULTY INSPECTION (U) MARYLAND UNIV COLLEGE PARK DEPT OF MANAGEMENT SCIENCES AND STATISTICS M SHISONG ET AL. JUN 85	1/1
UNCLASSIFIED	UMD-DM55-85/6 N00014-84-K-0301	F/G 12/1 NL

[illegible]



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963 A

ESTIMATION BY MOMENTS IN A MODEL OF FAULTY INSPECTION

Mao ShiSong  
East China Normal  
University, Shanghai

Samuel Kotz  
University of  
Maryland, College Park

Norman L. Johnson  
University of North  
Carolina, Chapel Hill

Key Words and Phrases: Attribute inspection; binomial distribution; factorial moments; hypergeometric distribution; mixtures; moment method.

ABSTRACT

Methodology developed by Blischke (Ann. Math. Statist. 33 (1962), 444-54) is applied to estimate the parameters in a model of faulty inspection, and to obtain approximate formulae for the variances of these estimators.

June 1985

310 FILE COPY

RECEIVED  
JUN 22 1985

## ESTIMATION BY MOMENTS IN A MODEL OF FAULTY INSPECTION

Mao ShiSong  
East China Normal  
University, Shanghai

Samuel Kotz  
University of Maryland  
College Park

Norman L. Johnson  
University of North  
Carolina, Chapel Hill

### 1. INTRODUCTION

Recent papers (Johnson et al. (1980), Johnson & Kotz (1983), Kotz & Johnson (1982)) have developed distributions of observed numbers of apparently defective items when sample inspection is imperfect, resulting in some defectives not being observed as such, and possibly some non-defective being described as 'defective' ("false positives"). Although these results are of interest, some more practical problems arise when it is desired to test whether the inspection is faulty or to estimate the degree of imperfection. In Johnson and Kotz (1985) some tests for detection of faulty inspection were investigated. The present paper is devoted to the estimation aspects of the problem. We will consider here the simplest form of inspection by attributes, assuming lot size to be, effectively, infinite. Each individual in a random sample of size  $n$  is examined and a decision reached as to whether or not it is 'nonconforming' (NC). Ideally, of course, such decisions should be completely free of error, but, as is well-known, this is often not the case. As a model of faulty inspection, we introduce two parameters

$$p = \text{Pr}[\text{individual declared NC} \mid \text{individual is NC}]$$

$$p' = \text{Pr}[\text{individual declared NC} \mid \text{individual is not NC}].$$

and suppose we wish to estimate these parameters. The proportion,  $P$ , of NC individuals in the lot is unknown, and plays the role, in this context, of a nuisance parameter.

A-1

It is clearly not possible to estimate  $p$  and  $p'$  (or  $P$ ) if each individual is examined only once. The only function of the parameters which can be estimated from such data is essentially  $Pp + (1-P)p'$  - the probability that an individual chosen at random is declared NC - because the distribution depends only on this quantity.

## 2. ESTIMATION

If individuals are examined more than once, however, it is possible to estimate each of the three parameters. We will suppose that each of the  $n$  individuals in the random sample is examined on  $m$  independent occasions. If  $D_i$  denote the number of times the  $i$ -th individual is declared to be NC, it has the distribution

$$\Pr[D_i = d_i] = P \binom{m}{d_i} p^{d_i} (1-p)^{m-d_i} + (1-P) \binom{m}{d_i} p'^{d_i} (1-p')^{m-d_i} \quad (1)$$

$$(d_i = 0, 1, \dots, m).$$

This is a mixture of two binomial distributions, with parameters  $(m, p)$  and  $(m, p')$  in proportions  $P$ ,  $(1-P)$  respectively. The  $r$ -th factorial moment of each  $D_i$  is

$$\begin{aligned} \mu_{(r)} &= E[D_i^{(r)}] \\ &= E[D_i(D_i-1) \dots (D_i-r+1)] = m^{(r)} \{Pp^r + (1-P)p'^r\} \end{aligned} \quad (2)$$

Estimating the parameters by making sample and population values of the first three factorial moments agree, we have

$$Pp^r + (1-P)p'^r = F_r \quad (r=1,2,3) \quad (3)$$

$$\text{where } F_r = (m^{(r)})^{-1} n^{-1} \sum_{i=1}^n D_i^{(r)}$$

Solutions  $\bar{p}$ ,  $\bar{p}'$  and  $\bar{P}$  of (3) are given by Jones (1933) as follows

(a)  $\bar{p}$ ,  $\bar{p}'$  are roots (in  $\theta$ ) of the equation

$$\theta^2 - A\theta + AF_1 - F_2 = 0 \quad (4)$$

where  $A = (F_3 - F_1 F_2) / (F_2 - F_1^2)$  (note that  $AF_1 - F_2 = (F_1 F_3 - F_2^2) / (F_2 - F_1^2)$ )

$$(b) \quad \bar{P} = (F_1 - \bar{p}') / (\bar{p} - \bar{p}') \quad (5)$$

There is indeterminacy in the solution, since if  $(\bar{p}, \bar{p}', \bar{P})$  is a solution, so is  $(\bar{p}', \bar{p}, 1 - \bar{P})$ . We will adopt the convention of regarding the greater root of (4) as the estimator ( $\bar{p}$ ) of  $p$ . It is reasonable to suppose that  $p$  is greater than  $p'$  - that is, the probability of declaring an individual to be NC if it is, indeed, NC is greater than if it is not. However, it must be remembered that even if this is so (i.e.  $p > p'$ ), this does not ensure that  $\bar{p}$  must exceed  $\bar{p}'$ .

### 3. Illustrative Example

For purposes of calculation, note that

$$\sum_{i=1}^n D_i^{(r)} = \sum_{j=1}^m N_j j^{(r)} \quad (6)$$

where  $N_j$  = number of individuals declared NC on just  $j$  occasions among the  $m$  times examined.

Suppose we have  $n=50$ ,  $m=3$ ;  $N_0=43$ ,  $N_1=1$ ,  $N_2=1$ ,  $N_3=5$ . Then

$$F_r = (3^{(r)} \cdot 50)^{-1} (43 \cdot 0^{(r)} + 1 \cdot 1^{(r)} + 1 \cdot 2^{(r)} + 5 \cdot 3^{(r)}) ,$$

is  $F_1 = \frac{1}{150} (1+2+15) = \frac{3}{25} ;$

$$F_2 = \frac{1}{300} (2+30) = \frac{8}{75} ;$$

$$F_3 = \frac{1}{300} \cdot 30 = \frac{1}{10} ,$$

where  $A = \frac{327}{346} = 0.94508671$  and  $\bar{p}, \bar{p}'$  are roots of  $\theta^2 - 0.94519\theta + 0.00674 = 0$ .

We find  $\bar{p} = 0.9379$ ;  $\bar{p}' = 0.0072$ ;  $\tilde{P} = 0.1212$ .

In this case  $\bar{p}, \bar{p}'$  and  $\tilde{P}$  are all between 0 and 1.

If they are not the method fails, though Blischke (1962, 1964) has suggested rules for this case.

#### 4. Variances

Blischke (1962) has obtained the following asymptotic formulae for the variances of  $\bar{p}, \bar{p}'$  and  $\tilde{P}$ :

$$\text{var}(\bar{p}) = \frac{p(1-p)}{P \cdot nm} + \frac{2(4B_2 + B_2')}{nm(m-1)C_2} + \frac{6(B_3 + B_3')}{nm(m-1)(m-2)C_4} \quad (7.1)$$

$$\text{var}(\bar{p}') = \frac{p'(1-p')}{(1-P)nm} + \frac{2(B_2 + 4B_2')}{nm(m-1)C_2'} + \frac{6(B_3 + B_3')}{nm(m-1)(m-2)C_4'} \quad (7.2)$$

$$\text{var}(\tilde{P}) = \frac{P(1-P)}{n} + \frac{18(B_2 + B_2')}{nm(m-1)(p-p')^4} + \frac{24(B_3 + B_3')}{nm(m-1)(m-2)(p-p')^6} \quad (7.3)$$

where  $B_h = Pp^h(1-p)^h$ ;  $B'_h = (1-P)p'^h(1-p')^h$

$$C_h = P^2(p-p')^h; C'_h = (1-P)^2(p-p')^h$$

When  $m$  is large, the first term of each expression usually gives quite good approximation, so that we may take

$$\text{var}(\tilde{p}) \div \frac{p(p-1)}{Pmn}; \text{var}(\tilde{p}') \div \frac{p'(1-p')}{(1-P)mn}; \text{var}(\tilde{P}) \div \frac{P(1-P)}{n}. \quad (8)$$

Note that the denominators are, for  $\text{var}(\tilde{p})$ , the expected number of examinations of NC individuals; for  $\text{var}(\tilde{p}')$ , the expected numbers of examinations of conforming individuals; and for  $\text{var}(\tilde{P})$  the number of individuals in the sample.

Using the numerical values in the example of Section 3, and inserting the values  $\tilde{p}, \tilde{p}', \tilde{P}$  for  $p, p', P$  respectively, we find (using (7.1)-(7.3)

$$\text{var}(\tilde{p}) \div 0.004133$$

$$\text{var}(\tilde{p}') \div 0.0000591$$

$$\text{var}(\tilde{P}) \div 0.002170$$

The last two terms in the expressions on the right-hand sides of (7.1)-(7.3) are

$$0.000885 \text{ and } 0.000044 \text{ for } \text{var}(\tilde{p});$$

$$0.000004 \text{ and } 0.000001 \text{ for } \text{var}(\tilde{p}');$$

$$0.000036 \text{ and } 0.000003 \text{ for } \text{var}(\tilde{P}).$$

So the use of (8) in this case, at least, would give quite good results,



even though  $m$  is only 3. (To the same order of approximation the three estimators are uncorrelated.)

### 5. Confidence Intervals

Blischke (1962) also showed that the asymptotic distributions of the estimators are normal. For  $P$ , approximate  $100(1-\alpha)\%$  confidence regions can be obtained from the inequality

$$\frac{n(\bar{P}-P)^2}{P(1-P)} < \lambda_{\frac{1}{2}\alpha}^2$$

where  $\Phi(-\lambda_{\frac{1}{2}\alpha}) = \frac{1}{2}\alpha$  and  $\Phi(y) = (\sqrt{2\pi})^{-1} \int_{-\infty}^y e^{-\frac{1}{2}u^2} du$ . Taking  $\alpha = 0.05$ , so that  $\lambda_{\frac{1}{2}\alpha}^2 = \lambda_{0.025}^2 = 3.8416$  the approximate 95% region for  $P$  is

$$50(0.1212-P)^2 < 3.8416 P(1-P)$$

or equivalently

$$53.8416 P^2 - 15.9616P + 0.73447 < 0$$

that is  $0.057 < P < 0.240$ .

Unfortunately we cannot use this method to obtain confidence regions for  $p$  and  $p'$ . The corresponding region for  $p$  (using (8)) would be

$$\frac{P mn(\bar{p}-p)^2}{p(1-p)} < \lambda_{\frac{1}{2}\alpha}^2$$

which cannot be used because  $P$  is not known. We might replace  $P$  by  $\bar{P}$ . This would give an asymptotically correct region.

Using the value  $\bar{P} = 0.1212$ , and taking  $\alpha = 0.05$ , as before, we get the (approximate) 95% confidence regions:

for  $p$ :  $0.1212 \cdot 150(0.9379-p)^2 < 3.8416 p(1-p)$  ,

whence  $22.0216 p^2 - 37.9436p + 15.9922 < 0$

leading to the interval  $0.7350 < p < 0.9880$  ;

for  $p'$ :  $18.18 (0.0072-p')^2 < 3.8416 p'(1-p')$

whence  $22.0216 p'^2 - 4.1034 p' + 0.00094 < 0$

leading to the interval  $0.00023 < p' < 0.0650$ .

### References

Blischke, W.R. (1962) Moment estimators for the parameters of a mixture of two binomial distributions, Ann. Math. Statist., 33, 444-454.

Blischke, W.R. (1964) Estimating the parameters of mixtures of binomial distributions, J. Amer. Statist. Assoc., 59, 510-528.

Johnson, N.L. and Kotz, S. (1983) "Faulty Inspection Distributions - Some Generalizations". In Reliability in the Acquisitions Process (D.J. DePriest and R.L. Launer, Eds.) pp. 171-182. M. Dekker, New York.

Johnson, N.L. and Kotz, S. (1985) Some tests for detection of faulty inspection, Statist. Hefte, 26, 19-29.

Johnson, N.L., Kotz, S. and Sorkin, H.L. (1980) Faulty inspection distributions, Commun. Statist. - Theory Methods, 9, 917-922.

Jones, H.G. (1933) A note on the n-ages method. J. Inst. Actu., 64, 318-324.

Kotz, S. and Johnson, N.L. (1982) Errors in inspection and grading: Distributional aspects of screening and hierarchical screening. Commun. Statist. Theory Methods, 11, 1997-2010.

### ACKNOWLEDGMENTS

Dr. Samuel Kotz's work was supported by the U.S. Office of Naval Research under Contract N 00014-84-K-301.

Professor Mao Shisong's work was carried out during a visit to the University of Maryland, College Park in 1985.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UMD DMSS-85/6			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Dep't. of Mgmt. & Stat.		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) University of Maryland College Park, Md. 20742			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION U.S. Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER ONR N00014-84-k-0301		
8c. ADDRESS (City, State, and ZIP Code) Arlington, Va. 22217			10. SOURCE OF FUNDING NUMBERS		
PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.		
11. TITLE (Include Security Classification) Estimation of Moments in a Model of Faulty Inspection					
12. PERSONAL AUTHOR(S) Mao ShiSong, Samuel Kotz and Norman L. Johnson					
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 3/85 TO 6/85		14. DATE OF REPORT (Year, Month, Day) June 1985	
15. PAGE COUNT (8) eight					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Attribute inspection; Binomial distribution; Mixtures; Moments method; Factorial moments.		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Methodology developed by Blischke (Ann. Math. Statist. 33)(1962), 444-54) is applied to estimate the parameters in a model of faulty inspection, and to obtain approximate formula for the variances of these estimators.					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Samuel Kotz			22b. TELEPHONE (Include Area Code) 301- 454 6805		22c. OFFICE SYMBOL

**END**

**FILMED**

**8-85**

**DTIC**